


bioNexia® Strep A dipstick is a rapid test

for the qualitative detection of group A Streptococcal antigen (GAS)


directly from human throat swab specimens



TEST PROCEDURE









Why choose

bioNexia® Strep A dipstick?

- Aid in the determination of the presence of group A *Streptococcus*
- Result available in just 5 minutes enabling physicians to make a quick diagnosis and prescribe suitable treatment
- Result remains stable for an additional 5 minutes
- Affordable strip format
- Easy-to-use, all-inclusive kit



Clinical Background

- Acute sore throat is a symptom often caused by an inflammatory process in the pharynx, tonsils or nasopharynx. Most of these cases are of viral origin and occur as part of the common cold,¹ requiring only symptomatic treatment. However, when bacterial pathogens are the cause of pharyngeal infections, antibiotic treatment may be required.
- Beta-hemolytic group A Streptococcus (GAS) is a major cause of upper respiratory tract infection, such as tonsillitis or pharyngitis. Rapid, accurate diagnosis of a GAS infection can help physicians to prescribe antibiotic treatment only when necessary, helping to reduce incidence of adverse effects and the spread of bacterial resistance.
- Early diagnosis and treatment of group A streptococcal pharyngitis has also been shown to reduce the severity of symptoms and further complications, such as rheumatic fever and glomerulonephritis.²,³
- 1. Pelucci, C., Grigoryan, L., Galeone, C. Guideline for the management of acute sore throat. ESCMID Sore Throat Guideline Group. Clin Microbiol Infect. 2012;18:1-27.

 2. Facklam, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, P. Marchiologies, Societies, Microbiology, 100. P.DISCA, 178.

 2. Facklam, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, H.J.

 2. Galden, R.R. and Carey, R.B., Streptococci and Aerococci, Manual of Clinical Microbiology, 4th ed., Lennette, E.H., Balows, A., Hausler, W.J. and Shadomy, A. C. and R. and R.
- editors, American Society for Microbiology, 1985, P154-175.

 3. Edwards, E.A., Phillips, I.A., and Suiter, W.C. Diagnosis of group A streptococcal infections directly from throat secretions, J. Clin. Microbiol. 1982;15:481-483.

Technical data:

- Reference: 414 556 (25 tests)
- Analytical sensitivity: 1x10⁴ org/swab
- Shelf-life: 24 months from date of manufacturing
- Storage temperature: +2°C to +30°C
- Content of the kit
 - 25 test strips in individually sealed pouches
- 25 sterile swabs
- 25 extraction tubes
- 25 sterile tongue depressors
- Two reagents: R1 and R2
- Controls: 1 positive, 1 negative
- 1 quick user guide
- 1 package insert
- 1 reagent holder

Dimensions of the kit

- Length: 220 mm
- Width: 180 mm
- Height: 86 mm
- Weight: 432 g

Want MORE from bioNexia?

Explore More Products, More Features and More Services at www.biomerieux.com/bionexia







bioMérieux S.A. 69280 Marcy l'Etoile **France**

Tel.: +33 (0)4 78 87 20 00 Fax: +33 (0)4 78 87 20 90

